

Temperature dependence of phenanthrene cavity radius in apolar solvents and in water

I. A. Ar'ev¹ and N. I. Lebovka²

¹ *Institute of Colloid and Water Chemistry named after A. V. Dumanskii, NASU, bulv. Vernadskogo, 42, 03142, Kyiv, Ukraine*

² *Biocolloid Chemistry Institute named after F. D. Ovcharenko, NASU, bulv. Vernadskogo, 42, 03142, Kyiv, Ukraine*

Phenanthrene, a three ring aromatic hydrocarbon, is used as a model substance for study of intermolecular interactions in dilute solutions of organic solvents and water by spectroscopic method. Temperature dependencies of shift of the $S_2 \leftarrow S_0$ spectrum of phenanthrene dissolved in apolar solvents and in liquid water are studied in this work. The spectroscopic data are used for analysis of the cavity radius R for phenanthrene molecule in solvents. It is shown that the value of R increases with temperature in organic solvents. In contrast, for water solution, the value of R initially grows with temperature increase (in the interval of 273.3-296 K), but becomes constant at higher temperatures. The changes of water structure in the neighbourhood of a phenanthrene molecule and probable hypothesis about causes of R value constancy in high-temperature region are discussed. Conclusion about strengthening of the water structure in an aqueous cage with phenanthrene molecule inside is done.

PACS numbers: 64.75.+g; 82.30.Rs; 61.20.Qg; 61.25.Em

Keywords: spectral shift, temperature expansion, hydrophobic hydration.

I. INTRODUCTION

Solute-solvent distribution function is the main characteristic for description of the structure of different molecular solutions at molecular level. A rather popular problem in this field is one related to study of the different aspects of hydrophobic hydration. Hydrophobic effects control solubility of apolar molecules in an aqueous solvents, formation of micelles, protein folding, etc. [1]. Hydrophobic hydration is a driving force of the hydrophobic effect. Unusual distribution of water molecules around an apolar solute or an apolar fragment of biphyllic molecule is the unique property of hydrophobic hydration. In spite of sustained efforts undertaken during last decades and existence of a lot of theoretical speculations, this matter is still poorly studied experimentally.

This situation is caused by the lack of effective empirical tools for structural studies in the nearest vicinity of hydrophobic molecules dissolved in water. The aqueous solubility of hydrophobic molecules in the majority of cases are very low. Traditional experimental methods, e.g. neutrons or x-rays scattering, are ineffective in application to such diluted solutions.

The same difficulty is encountered also for nonaqueous diluted solutions, where both solute and solvent molecules are apolar. Solutions of apolar compounds in apolar solvents can serve as reference systems. For these solutions the contribution of electrostatic interactions can be neglected.

The spectral shift method [2] can be effective in such a situation, though it is somewhat rough. As a rule, transfers of a molecule from one environment to another one induces shifts in its electronic spectrum. The spectral shift $\Delta\nu$ depends on molecular environment of the test molecule. The method of spectral shifts can be used for estimation of the mean distance between the solute and solvent molecules when both solute and solvent molecules are apolar. This distance may be identified with a certain approximation as $R + r$, where R is the radius of the cavity, which contains a solute molecule, and r is the radius of the solvent molecule. Note that the first maximum of the solute-solvent pair distribution function is approximately localized at $R + r$. This method can be applied also for polar solvents, though with definite restriction: the permanent electric fields induced on the solute by solvent molecules should be compensated.

Here we present results of investigation of the shift dependencies in $S_2 \leftarrow S_0$ spectrum $\Delta\nu$ of phenanthrene dissolved in some apolar solvents, in methanol and in water in the wide temperature ranges. Our purpose is to examine changes of R with temperature. The paper is organized as follows. In Sec. II we give introductory information about the solvent-induced spectral shift method. In Sec. III the experimental procedures are described. Experimental results are presented and discussed in Sec. IV. Conclusions are summarized in Sec. V

II. THE SPECTRAL SHIFT METHOD

Let an apolar molecule be transferred from a rarified gas into the liquid solvent. Then, in approximation of the dielectric continuum, the spectral shift is

$$\Delta\nu = -C\phi(R)f(n) + \Delta\nu_o \quad (1)$$

where, C is a positive factor that characterizes the solute molecule and $f(n) = (n^2 - 1)/(n^2 + 2)$ is the Lorentz-Lorenz function, n is the refractive index of the solvent, $\Delta\nu_o$ is related to contributions other than dispersion interactions. The $\phi(R)$ is a function of the cavity radius. When the solute molecule is located in the cavity center, $\phi(R) = 1/R^3$. When the solute molecule is in contact with the cavity surface, $\phi(R) = R^3/(r_u^3(2R - r_u)^3)$, where r_u is the solute radius.

Eq. 1 is good for approximation only in a case when the solute oscillator strength lies in the interval from 0.1 to 1 (see [2] and references cited therein).

Eq. 1 can be rewritten as

$$\phi(R) = (\Delta\nu_o - \Delta\nu)/(Cf(n)). \quad (2)$$

This equation may be used for determination of the temperature dependence of $R(T)$. In particular, in a case when the solute molecule is situated in the cavity center, it follows from Eq. 2 that

$$\frac{R(T)}{R(T_{ref})} = \left[\frac{(\Delta\nu_{0,ref} - \Delta\nu_{ref}) f(n)}{(\Delta\nu_0 - \Delta\nu) f(n_{ref})} \right]^{1/3}, \quad (3)$$

where subscript *ref* shows that the property was measured at some reference temperature T_{ref} .

The temperature coefficient of the spectral shift may be defined as

$$\alpha_s = (1/\Delta\nu)(d\Delta\nu/dT). \quad (4)$$

Then, in the case when $\Delta\nu_o$ contribution to Eq. 1 may be neglected (see [2] for more detailed discussion), and the solute is located in the cavity center, we obtain

$$\alpha_s = -\alpha_v - \alpha_c, \quad (5)$$

where α_v is the volume expansion temperature coefficient of the bulk solvent, and $\alpha = (3/R)(dR/dT)$ is the volume expansion temperature coefficient of the cavity, occupied by a solute. This equation determines the lower limit of α_s .

III. EXPERIMENTAL

All absorption spectra were recorded by spectrophotometer Specord UV Vis using the following procedure. First, the reference spectrum of lanthanide salts dissolved in fused quartz was recorded, then the spectrum of the studied sample was recorded and then the reference spectrum was recorded again. The registration scale was $20.2 \text{ cm}^{-1}/\text{mm}$, the registration rate was $500 \text{ cm}^{-1}/\text{min}$. The spectrograms were suitable for the next treatment if the distance between the band maximums in the first and the last reference spectrograms did not exceed 2 cm^{-1} . The wave number of the phenanthrene band maximum was measured for each spectrogram and averaged for all the data. The root-mean-square errors (rms) of wave-number positions did not exceed 2 cm^{-1} .

We used the twice-distilled water, chemically pure Phenanthrene and pure organic solvents for chromatography. The concentration of phenanthrene was of the order of 10^{-6} mol/dm^3 for aqueous solution (5 cm cell) and of the order of $10^{-4} - 10^{-5} \text{ mol/dm}^3$ (1 - 4 mm cells) for other solutions. Cylindrical cells and specially designed temperature chamber were used. Water was serving as a heat-carrier. Temperature of solutions was stabilized with a precision of 0.1 K.

Temperature shifts of phenanthrene vapor spectrum were measured using a procedure, different from that applied for solutions. The pressure of phenanthrene vapor in the preliminary pumped out and soldered 2-cm quartz cell was of the order of 10^{-2} mm Hg . The sample was heated in the special chamber to the temperatures from 360 to 615 K. The temperature variations did not exceed 0.2 K.




FIG. 1: The $S_2 \leftarrow S_0$ spectral shift, $\Delta\nu$, of phenanthrene versus Lorentz-Lorenz function, $f(n) = (n^2 - 1)/(n^2 + 2)$, in dilute solutions of n-pentane (C5), n-heptane (C7), n-decane (C10), n-hexadecane (C16), benzene, and water. The temperature T in K is indicated near the experimental points. The more large scale presentation for water is given at Fig. 2.

The temperature dependence of the long-wave band in $S_2 \leftarrow S_0$ Phenanthrene vapor spectrum was approximated by the linear equation

$$\nu = \nu_o - k_\nu T, \quad (6)$$

where $\nu_o = 35654 \pm 18 \text{ m}^{-1}$, $k_\nu = 1.175 \pm 0.038 \text{ m}^{-1}\text{K}^{-1}$, is the absolute temperature. The correlation coefficient of rms approximation was 0.999. Eq. 6 is qualitatively in agreement with analysis data for the $S_1 \leftarrow S_0$ temperature shift of anthracene transition in the gas phase [3]. So, we can conclude that this transition is of the same nature as $S_2 \leftarrow S_0$ transition of phenanthrene [4].

All necessary reference data were taken from [5, 6, 7, 8, 9].

IV. RESULTS AND DISCUSSION

Figure 1 shows experimental data for the spectral shifts $\Delta\nu$ of phenanthrene versus Lorentz-Lorenz function $f(n) = (n^2 - 1)/(n^2 + 2)$ in dilute solutions of organic solvents, and water at different temperatures.

Table 1 shows values of α_v , α_s , $-\alpha_v - \alpha_s$, for phenanthrene solutions at $T=293 \text{ K}$. Data for anthracene solutions are included for comparison (were taken from [10] or re-measured).

Although the anthracene polarizability exceeds that of phenanthrene, the observed values of α_s and $\alpha_c = -\alpha_v - \alpha_s$ are of the same order of magnitude within the experimental errors. It is surprising at first glance, because the more is polarizability, the more is intermolecular attraction interactions. So, values of α_s and α_c for these substances should be noticeably different. But phenanthrene molecule is more compact than anthracene one. So the cavity size for this molecule is more small and the attraction interactions for the phenanthrene are more pronounced than for the anthracene.

Table 1. Solvent thermal expansion coefficients (α_v), shift temperature coefficients (α_s), and $-\alpha_v - \alpha_s$ at 293 K for




FIG. 2: The $S_2 \leftarrow S_0$ spectral shift, $\Delta\nu$, of phenanthrene versus Lorentz-Lorenz function, $f(n) = (n^2 - 1)/(n^2 + 2)$, in dilute water solution. The temperature T in K is indicated near the experimental points.

phenanthrene (Ph) and anthracene (A). The temperature ranges used for determination of α_s values are shown in parentheses.

Solvent	$\alpha_v, 10^{-3}, \text{K}^{-1}$	$\alpha_s, 10^{-3} \text{K}^{-1}$		$-\alpha_s - \alpha_v, 10^{-3} \text{K}^{-1}$	
		Ph	A	Ph	A
Benzene	1.20	-1.69 (279.9-341.1 K)	-1.70 (283-333 K)	0.49	0.50
n-Heptane	1.13	-1.94 (277.6- 360.1 K)	-1.92 (274.2-309.1 K)	0.81	0.79
n-Decane	1.05	-1.75 (282.4-338.2 K)	-1.75 (275.4-353.1 K)	0.70	0.70
n-Hexadecane	0.88	-1.49 (293.1 - 355.6 K)	-	0.61	-
Methanol	1.14	-1.63 (276.4-323.3 K)	-1.64 (293.1 - 324.3 K)	0.48	0.50
Acetone	1.41	-	-2.25 (276.6-321.7, K)	-	0.84

The values of $-\alpha_s - \alpha_v$ represent temperature expansion of cavities in apolar solvents. These values for polar solvents are also non-zero and positive.

Data for water spectral shift $\Delta\nu$ in the temperature range from 296 to 363.6 K the may be approximated by linear equation 1. The equation parameters are $C\phi(R) = 6250 \pm 50 \text{ cm}^{-1}$, $\Delta\nu_0 = 97 \pm 10 \text{ cm}^{-1}$, and correlation coefficient is 0.998 (dashed line in Fig. 2). It should be noted that these parameters were estimated proceeding from an assumption that terms of Eq. 6 are $\nu_o = 35654 \text{ m}^{-1}$ and $k_\nu = 1.175 \text{ m}^{-1} \text{K}^{-1}$ exactly. Taking into account errors of estimations, one cannot state that $\Delta\nu_0 \approx 0$ and that means that electrostatic fields on phenanthrene can be canceled effectively. At low temperatures (in the interval of 273.3-296 K) the pronounced deviations from linear relation is observed.

Figure 3 presents the temperature dependencies of relative phenanthrene cavity radius $R(T)/R_{300}$ in organic solvents and water estimated from Eq. 3. Here R_{300} corresponds to the cavity radius at reference temperature $T = 300 \text{ K}$.

The cavity radius continuously grows with temperature in organic solvents. It is interesting to note that R value of grows more noticeably within congeneric series $\text{C}_7 - \text{C}_{16}$ than in benzene solutions. In contrast the R value of water solution initially grows with temperature increase (in the range of 273.3-296 K), but becomes constant at higher temperatures. The explanation of this experimental observation is not trivial.

This fact can evidence higher structurization of the aqueous cage with phenanthrene molecule inside at high temperatures. It is in accordance with conclusion made in [11] regarding strengthening of the hydrogen bonds in the first




FIG. 3: Temperature dependencies of relative phenanthrene cavity radius $R(T)/R_{300}$ in n-heptane (C_7), n-decane (C_{10}), n-hexadecane (C_{16}), benzene and water.

hydration shell of a solute molecule and formation of a more organized structure around the solute. Tanaka [12] has also observed in his simulations the effect of structural enhancement in water near a large cavity.

It is possible that at temperatures exceeding 296 K, the H-bonds between water molecules get strengthened in the nearest vicinity of phenanthrene to such extent that thermal factor not affect them. Tomchuk and Krasnopolovets [13] have shown that if the chain of water molecules is not perturbed thermally, then all the protons in the chain vibrate coherently together jumping from one O - H bond to the other one. Two ions of the opposite signs arise on the ends of such chain as a result of their first jump, the next jump restores the initial position and so on. This process has a tunneling nature [13]. Note that the number of O - H bonds decreases by one during the first jump. Hence, the total energy of the chain should increase by the energy of one H-bond breakage. The loss of the bond should be compensated by electronic polarization that results in the strengthening of the hydrogen bonds. But the energy of the chain does not increase. The molecular beams experiments [14] support this assumption. The electronic polarization of water molecules can also play an important role in process of electron tunneling through the water layers [15]. So, the coherent proton vibration can promote stabilization of the chains of water molecules.

V. CONCLUSIONS

We have measured temperature shifts of $S_2 \leftarrow S_0$ in the spectra of gaseous phenanthrene and phenanthrene dissolved in some organic apolar and polar solvents and in water. The temperature expansions of cavities containing solute molecules inside, α_c 's, were calculated for the apolar solvents as differences between temperature coefficient of the shift, α_s , and temperature expansion of the solvent, α_v , and were compared with the same values for anthracene solutions. It was shown that the value of R for phenanthrene increases with temperature in organic solvents. The aqueous system shows different properties at temperatures higher than 296 K and below this temperature. In contrast, the value of R in water solution of phenanthrene initially grows with temperature increase (in the range of 273.3-296 K), but becomes constant at higher temperatures. These phenomena were related to strengthening of the hydrogen bonds in the first hydration shell of a solute molecule and formation of a highly organized structure around the solute.

ACKNOWLEDGEMENTS

Authors thank Dr. N.S. Pivovarova for her help with preparation of the manuscript.

-
- [1] W. Blokzij and J. B. F. N. Engberts, "Hydrophobic effects: Opinions and facts", *Angew. Chem. Int. Ed. Engl.* **32**, pp. 1545 - 1579, 1993.
 - [2] I. A. Ar'ev, "Study of benzene hydrophobic hydration by the spectral shift method" *Teor. Eksp Khim.* **23**, pp 329-339, 1987.
 - [3] S. O. Mirumyants and Yu. S. Demchuk, "On the discovering of the parameters determining the broadening of multiatom-molecule vapour vibronic spectra". *Optika i spektr.* **56**, 241-246, 1984.
 - [4] E. Clar, "Polycyclic hydrocarbons" (Russian translation). Khimia, Moscow, 1971.
 - [5] B. V. Ioffe, "Refractometric methods in chemistry" (in Russian). Goskhimizdat, Moscow, 1960.
 - [6] I. B. Vargaftik. Handbook on thermophysical properties of gases and liquids (in Russian). Nauka, Moscow, 1972.
 - [7] Short handbook on physical and chemical values (in Russian). Ed. by A. A. Ravdel and A. M. Ponomaryova. 8-th edition remade. Khimiya, Leningrad, 1983.
 - [8] CRC Handbook of Chemistry and Physics. Editor-in-Chief R. C. West. 66th ed., Boca Raton, Florida: CRC Press, 1985 - 1986.
 - [9] I. T. Goronovskii, Yu. P. Nazarenko and E. F. Nekryach. Short handbook on chemistry (in Russian). Naukova dumka, Kiev, 1987.
 - [10] I. A. Ar'ev and N. I. Lebovka, "Methods of effective molecular size evaluation and temperature shifts of $S_1 \leftarrow S_0$ anthracene spectrum". *Proceedings of SPIE*, **4938**, pp. 105 - 112, 2002.
 - [11] S. Urahata and S. Canuto, "Monte Carlo study of the temperature dependence of the hydrophobic hydration of benzene". *Chem. Phys.Lett.*, **311**, pp. 235 - 240, 1999.
 - [12] H. Tanaka, "Cavity distribution in liquid water and hydrophobic hydration". *Chem. Phys. Lett.*, **282**, pp. 133 - 138, 1998.
 - [13] P. M. Tomchuk and V. V. Krasnolovets, "Macroscopic quantum tunneling of polarization in the hydrogen-bonded chain". *J. Molec. Struct.*, **416**, pp. 161 - 165, 1997.
 - [14] E. P. Grimsrud and P. Kebarle, "Gas phase ion equilibria studies of solvation of the hydrogen ion by methanol, dimethyl ether, and water. Effect of hydrogen bonding". *J. Amer. Chem. Soc.*, **95**, pp. 7939 - 7943, 1973.
 - [15] A. Mosyak, P. Graf, I. Benjamin, and A. Nitzan, "Electron tunneling through water layers: Effect of polarizability". *J. Phys. Chem. A*, **101**, pp. 429 - 433, 1997.